

Research Article

Knowledge, Attitude, and Practice toward Prevention of COVID-19 Among Nursing and Midwifery Students: A cross-sectional Study in Gorgan, Northern Iran

Iman Taghizadeh Firoozjaie¹, Homeira Khoddam^{2*}, Mohamadreza Mohamadi³, Mahin Tatari², and Seyedmahrokh A. Maddah⁴

¹Department of Medical Surgical Nursing, School of Nursing and Midwifery, Golestan University of Medical Sciences, Gorgan, Iran

²Nursing Research Center, Golestan University of Medical Sciences, Gorgan, Iran

³Department of Psychiatry, School of Medicine, Neuroscience Research Center, Golestan University of Medical Sciences, Gorgan, Iran

⁴Department of Anesthesiology, Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, Iran

ORCID:

Iman Taghizadeh Firoozjaie: <https://orcid.org/0000-0002-3150-4178>

Homeira Khoddam: <https://orcid.org/0000-0002-8225-2652>

Mohamadreza Mohamadi: <https://orcid.org/0000-0001-5525-222X>

Mahin Tatari: <https://orcid.org/0000-0003-0781-7139>

Seyedmahrokh A Maddah: <https://orcid.org/0000-0002-3605-6917>

Corresponding Author:

Homeira Khoddam; email:

drmahnazmodanloo@yahoo.com

Received 22 Jan 2022

Accepted 19 June 2022

Published 30 September 2022

Production and Hosting by
Knowledge E

© Firoozjaie et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editor-in-Chief:

Prof. Nazik Elmalaika Obaid
Seid Ahmed Husain, MD,
M.Sc, MHPE, PhD

Abstract

Background: Healthcare providers are high-risk groups for the novel coronavirus disease (COVID-19) infection. Nursing students are an essential part of high-risk healthcare providers. The aim of this study was to evaluate the knowledge, attitude, and practice of nursing and midwifery students at Golestan University of Medical Sciences (GOUMS) toward the prevention of COVID-19.

Methods: This cross-sectional study was conducted on 174 nursing and midwifery students of GOUMS (northeast of IRAN) in 2020. Data were gathered using an online questionnaire comprising three parts – Knowledge (8 questions), Attitude (10 questions), and Practice (8 questions) toward the prevention of COVID-19. The range of scores in each subscale (knowledge, attitude, and preventive practices) were 8–24, 10–50, and 8–24, respectively. Data were analyzed using the SPSS v.16 software. The Mann–Whitney and Kruskal–Wallis tests were used for the analysis.

Results: Overall, in this study, nursing and midwifery students had a good knowledge (mean score: 23.19 ± 2.56), a positive attitude (mean score: 45.48 ± 4.21), and appropriate practice (mean score: 23.30 ± 3.51) regarding COVID-19 prevention. Also, the results revealed a positive correlation between knowledge and attitude ($r = 0.1$, $P = 0.18$) and attitude and practice ($r = 0.2$, $P = 0.01$) among the students.

Conclusion: The findings demonstrated a good preventive knowledge, attitude, and practice toward COVID-19 among nursing and midwifery students.

Keywords: health knowledge, attitudes, practice, COVID-19, students, nursing, Iran

OPEN ACCESS

1. Introduction

The novel coronavirus disease (COVID-19) belongs to the Betacoronas family, which was first reported in Wuhan, China in 2019. COVID-19 is very contagious and, similar to its other siblings, can produce a wide spectrum of symptoms such as common cold or severe respiratory problems in MERS-COV or SARS-COV [1]. The COVID-19 virus had a high transmissibility rate and global rapid spread in a short period of time [2]. On June 2nd, 2020, the World Health Organization (WHO) reported a total of 376,320 deaths due to COVID-19 [3].

These viruses are Zoonotic in origin with some types transferred from cats and camels. Although there are several types of viruses in this family, no human involvement is seen till now [4]. This virus, with its surface protein (S) can bind to the host receptor angiotensin converting enzyme 2 (ACE2) to enter the host cell. ACE2 is a peptidase and is attached to the cell membrane in several tissues throughout the body, including the heart, gastrointestinal tract, kidneys, and lungs [5–7]. The most common symptoms of COVID-19 infection are respiratory symptoms, fever, cough, dyspnea, malaise, and fatigue. In severe forms it can lead to pneumonia, acute respiratory distress syndrome (ARDS), renal failure, and also death [8–10]. Acute hemorrhagic encephalopathy and increasing risk of cerebrovascular accidents (CVA) have also been seen and reported [11, 12]. In previous studies in China, mortality was 0.9% in patients without underlying medical conditions. Patients with underlying cardiac disease, diabetes mellitus (DM), and hypertension (HTN) were at higher risk of mortality [13].

Standard recommendations for preventing the infection such as handwashing, wearing masks, and maintaining social distancing are vital. However, social media being the main source of news and information today, most people are also exposed to inaccurate and fake information online that can develop or amplify fear and anxiety among them [9, 14]. However, based on research evidence, patients felt most assured when healthcare providers were the source of information [2].

The knowledge, attitude, and practice of people plays a fundamental role in controlling a disease/pandemic [15]. Lack of information about can lead to an ultimate disaster [16]. For assurance of practical preventive actions and optimal readiness, providing adequate information, educational packages, and hygienic strategies to society is vital in controlling system [17]. Considering nursing students are at the forefront of the healthcare today, they must have adequate knowledge and training for the prevention and appropriate management of COVID-19. Therefore, of the present study aims to

evaluate the knowledge, attitude, and practice of Iranian nursing and midwifery students toward the prevention of COVID-19.

2. Materials and Methods

This cross-sectional study was conducted on nursing and midwifery students of the Golestan University of Medical Sciences (GOUMS, northeast IRAN) in 2020. Following the approval of the Institutional Ethics Committee and receiving written informed consent from all study participants, 174 eligible students were recruited in the using census method. The inclusion criteria were being a nursing or midwifery student and active employment during study period. Cases with incomplete questionnaire were excluded. Data were collected using demographic information form and a “knowledge, attitude, and practice for prevention of respiratory tract infections” questionnaire which was developed by Goni *et al.* in 2020. Based on the viewpoint of experts, the questionnaire scoring was revised. Relying on the insight gathered from the expert panel, responses to questions were changed in each domain and then a pilot study was performed on 20 students from the same study population. The result of Cronbach’s alpha (0.85) indicated that the research instrument was reliable [18]. This self-administered questionnaire consisted of three domains – knowledge (8 items), attitude (10 items), and practice (8 items).

The knowledge domain of questionnaire consisted of eight items to assess students’ knowledge about the etiology, virus dissemination and disease transmission, signs and symptoms, complications, risk factors, and some protective methods for preventing of disease. Scoring ranged from 1 to 3 for wrong answers, don’t know, and correct answers, respectively. The total score in this part ranged from 8 to 24.

The second domain of the checklist evaluated students’ attitude toward the prevention of infection. This checklist had 10 questions which were scored on a five-point scale (1 = never; 2 = rarely; 3 = sometimes; 4 = often; 5 = always) and the total score in this part ranged from 10 to 50.

The third domain comprised eight questions which were scored on a three-point scale (always = 3, occasional = 2, never = 1), the maximum score was 24 and the minimum was 8.

For data collection, first an electronic questionnaire was designed on a virtual e-learning system. An online questionnaire link was sent to the students. Data were analyzed using the SPSS v.16 software (SPSS Inc., Chicago, IL, USA). Kolmogorov–Smirnov test was used to examine the normality of the variables, which showed a non-normal

distribution. Mann–Whitney test and Kruskal–Wallis test were used to determine the relationship between demographic information with knowledge, attitude, and practice of students. Also, Spearman's correlation test was used to measure whether and how much the variables are correlated [18].

3. Results

The mean age of the students was 21.97 ± 2.72 years (range, 18–38). Most students were nursing students (86.8%), single (89.1%), female (60.9%), and living in the city (83.3%). The biggest source of information was virtual information (35.1%). Only 4.6% of students were infected with COVID-19 before study, and 9.2% had family members infected with COVID-19. The underlying diseases were present in 17.2% of students and 31.6% of students had history of influenza vaccination before the study (Table 1).

The mean and standard deviation of scores of three domains were 23.19 ± 2.56 , 45.48 ± 4.21 , and 23.30 ± 3.51 , respectively.

There was no significant correlation between the mean scores of knowledge with demographic characteristics, source of information, and history of participants' infection by COVID-19 and history of their family members' infection and gaining awareness about the disease (Table 2).

The results of this study showed that the mean scores of attitude had no correlation with the knowledge, demographic characteristics, source of information, and COVID-19 infection in students or their family (Table 2).

The mean practice scores in students had a significant correlation with gender, marital status, and history of previous infection with COVID-19 in students and their family members (Table 2).

Moreover, the results showed no significant correlation between knowledge and attitude ($r = 0.1$, $P = 0.18$). There was an inverse correlation between students' knowledge and practice, however, this correlation was not statistically significant ($r = -0.05$, $P = 0.51$). Nevertheless, there was a positive correlation between students' attitude and practice ($r = 0.2$, $P = 0.01$).

4. Discussion

This study showed that nursing and midwifery students had a good and appropriate level of knowledge, attitude, and practice towards the prevention of COVID-19. The

TABLE 1: Frequency distribution of demographic and clinical characteristics of students.

Percentage	Frequency	Variable	
39.1	68	Male	Gender
60.9	106	Female	
89.1	155	Single	Marital status
10.9	19	Married	
83.3	145	City	Place of residence
15.5	27	Village	
86.8	151	Nursing	Field of study
13.2	23	Midwifery	
9.2	16	TV	Information source
35.1	61	Social media network	
2.3	4	Educational sources	
6	1	Public information	
47/7	83	More than one sources	
5/2	9	Others	
95.4	166	No	History of COVID-19 infection
4.6	8	Yes	
1.7	3	DM	Comorbid diseases
0.6	1	Cardiac disease	
3.4	6	HTN	
1.7	3	Immune deficiency disease	
0.6	1	COPD	
8	14	Asthma	
0.6	1	Allergic rhinitis	
82.8	144	None	
90.8	158	No	History of COVID-19 infection in family members
9.2	16	Yes	
68.4	119	No	Influenza vaccination
31.6	55	Yes	

results of a study on Palestinian nursing students revealed that they had a negative attitude and moderate level of knowledge about COVID-19 [19]. The results of another study indicated a high level of knowledge about COVID-19 among Philippine nursing students. Higher year level of students had significant correlation with their knowledge. In addition, female students had a higher attitude score than their male counterparts [20]. Another study in Vietnam revealed a good preventive practice toward COVID-19

TABLE 2: The knowledge, attitude and practice for prevention of COVID-19 disease among students based on demographic and clinical characteristics.

Practice		Attitude		Knowledge		Variable	
P-value	Mean $\pm$ SD	P-value	Mean $\pm$ SD	P-value	Mean $\pm$ SD		
0.04	22.26 $\pm$ 4.15	0.85	45.94 $\pm$ 4.42	0.33	23.40 $\pm$ 2.58	Male	Gender
	23.63 $\pm$ 3.30		46.00 $\pm$ 4.10		23.06 $\pm$ 2.55	Female	
0.003	23.40 $\pm$ 3.50	0.505	46.02 $\pm$ 4.34	0.93	23.17 $\pm$ 2.50	Single	Marital status
	20.57 $\pm$ 3.80		45.63 $\pm$ 3.04		23.36 $\pm$ 2.29	Married	
0.088	23.32 $\pm$ 3.68	0.3	46.04 $\pm$ 4.20	0.6	23.21 $\pm$ 2.56	City	Place of residence
	21.92 $\pm$ 3.8		45.41 $\pm$ 3.97		22.93 $\pm$ 2.63	Village	
0.078	22.90 $\pm$ 3.79	0.88	46.00 $\pm$ 4.26	0.37	23.28 $\pm$ 2.50	Nursing	Field of study
	23.34 $\pm$ 2.83		45.86 $\pm$ 3.95		22.60 $\pm$ 2.92	Midwifery	
0.622	23.40 $\pm$ 2.26	0.503	4.23 $\pm$ 2.26	0.577	24.00 $\pm$ 1.37	TV	Information source
	22.56 $\pm$ 3.74		45.13 $\pm$ 3.76		22.93 $\pm$ 2.74	Social media network	
	23.16 $\pm$ 3.55		45.47 $\pm$ 4.62		22.50 $\pm$ 2.52	Educational sources	
	23.75 $\pm$ 2.36		45.25 $\pm$ 7.23		23.00 $\pm$ 00	Public information	
	15.00 $\pm$ 00		44.00 $\pm$ 00		23.13 $\pm$ 2.59	More than one sources	
	23.27 $\pm$ 3.72		46.86 $\pm$ 3.77		24.00 $\pm$ 2.59	Others	
0.147	21.66 $\pm$ 4.09	0.81	43.33 $\pm$ 3.53	0.346	23.24 $\pm$ 2.52	Yes	History of COVID-19 infection
	23.19 $\pm$ 3.65		45.99 $\pm$ 4.25		22.00 $\pm$ 3.16	No	
	21.12 $\pm$ 4.58	0.477	45.75 $\pm$ 3.53	0.125	22.33 $\pm$ 3.05	DM	Comorbid diseases
	21.00 $\pm$ 2.64		44.66 $\pm$ 3.05		23.00 $\pm$ 00	Cardiac disease	
0.024	23.00 $\pm$ 00		44.00 $\pm$ 00		23.83 $\pm$ 2.04	HTN	
	23.83 $\pm$ 3.25		44.50 $\pm$ 3.88		23.33 $\pm$ 1.15	Impaired immune system	
	23.66 $\pm$ 2.30		47.33 $\pm$ 5.03		23.00 $\pm$ 00	COPD	
	23.00 $\pm$ 00		45.00 $\pm$ 00		23.21 $\pm$ 2.63	Asthma	
	20.00 $\pm$ 2.03		46.14 $\pm$ 3.73		17.00 $\pm$ 00	Allergic rhinitis	
	16.00 $\pm$ 00		38.00 $\pm$ 00		17.00 $\pm$ 00	Nothing	
0.009	23.36 $\pm$ 3.72	0.784	46.06 $\pm$ 4.26	0.121	23.11 $\pm$ 2.54	Yes	History of COVID-19 infection in family members
	23.32 $\pm$ 3.6		45.91 $\pm$ 4.26		23.94 $\pm$ 2.77	No	
0.519	20.87 $\pm$ 3.24	0.75	46.63 $\pm$ 3.84	0.819	23.21 $\pm$ 2.51	Yes	Influenza vaccination
	23.20 $\pm$ 3.62		45.87 $\pm$ 4.27		23.14 $\pm$ 2.68	No	

among healthcare students; however, sufficient knowledge and positive attitude were lower [21].

Goni *et al.*'s study in Malaysia revealed an appropriate and good level of knowledge for the prevention of influenza among Haj pilgrims [16]. However, other studies on Haj pilgrims showed low level of knowledge in this regard [22, 23]. In the present

study, participants had an excellent and appropriate level of knowledge. This difference could be due to the students' level of education. No significant correlation was seen between demographic variables and students' level of knowledge. Other studies did not confirm any correlation between demographic characteristics and participants' level of knowledge regarding viral infection prevention [24, 25]. This difference in correlation could be due to the study population and the difference in their cultures, age, and educational backgrounds.

In this study, the mean score in attitude toward coronavirus prevention was high. Previous studies revealed that participants had a positive attitude in observing preventive recommendations for MERS-CoV [26, 27]. The practice score in the prevention of COVID-19 had a significant correlation with the gender of participants. Females had more attention to prevention. Another factor which had a significant correlation was the history of comorbid diseases. One possible explanation for this is adequate education and knowledge of these participants and the high possibility of being infected by COVID-19 in patients with underlying disease, which is associated with higher knowledge and attitude [28]. History of previous infection of person or their family members with COVID-19 had a significant correlation with their preventive practice score. This could be due to students' previous personal experiences which can lead to better practice in this regard.

The results revealed a positive correlation between knowledge and attitude and attitude and practice among students. In an Indonesian study by Adli *et al.*, the level of knowledge of undergraduate medical students related to the COVID-19 pandemic was slightly positively correlated with their attitude and practice, while attitude was weakly correlated with their practice [29]. Another study in Ethiopia indicated a negative correlation between knowledge and attitude and between knowledge and practice regarding COVID-19 among healthcare workers [30].

5. Limitations

The present study has some limitations. Because of the COVID-19 pandemic, the students were mostly taking online classes through virtual and e-learning methods, resulting in limited access and interaction. The lack of generalizability of the findings to all nursing and midwifery student in different geo-cultural contexts, due to small sample size and participants' sociodemographic and cultural characteristics, is another limitations of this study..

6. Conclusion

The results of this study demonstrated a good preventive knowledge, attitude, and practice toward COVID-19 among nursing and midwifery students. However, additional education and training strategies are needed to offer a better understanding and motivate positive attitude and practice toward COVID-19 prevention for all nursing and midwifery students.

Acknowledgments

The authors are thankful to students for their participating in this study. Moreover, they are grateful to the administrators of the Faculty of Nurses and Midwifery for their support. They are also thankful to the Research Deputy of Golestan University of Medical Sciences for funding this study.

Ethical Considerations

This study was approved by institutional ethical research committee at Golestan University of Medical Science (Code: IR.GOUMS.REC.1399.006). All participants gave a written informed consent before filling in the questionnaire. Confidentiality of private information and voluntary participation was ensured.

Competing Interests

The authors have no conflicts of interest to declare for this study.

Availability of Data and Material

All relevant data of this study are available to any interested researchers upon reasonable request to the corresponding author.

Funding

This study was financially supported by the Deputy of Research and Technology, Golestan University of Medical Sciences, Gorgan, Iran.

References

- [1] Sharma, A., Ahmad Farouk, I., & Lal, S. K. (2021). COVID-19: A review on the novel coronavirus disease evolution, transmission, detection, control and prevention. *Viruses*, 13(2), 202. <https://doi.org/10.3390/v13020202>
- [2] Emami Zeydi, A., Ghazanfari, M. J., Sanandaj, F. S., Panahi, R., Mortazavi, H., Karimifar, K., Karkhah, S., & Osuji, J. (2021). Coronavirus disease 2019 (COVID-19): A literature review from a nursing perspective. *Biomedicine*, 11(3), 5–14. <https://doi.org/10.37796/2211-8039.1154>
- [3] WHO. *Coronavirus disease (COVID-2019) situation reports*. WHO. 2020.
- [4] Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G. F., Tan, W., & the China Novel Coronavirus Investigating and Research Team. (2020). A novel coronavirus from patients with pneumonia in China, 2019. *The New England Journal of Medicine*, 382(8), 727–733. <https://doi.org/10.1056/NEJMoa2001017>
- [5] Yaron, J. R., Ambadapadi, S., Zhang, L., Chavan, R. N., Tibbetts, S. A., Keinan, S., Varsani, A., Maldonado, J., Kraberger, S., Tafoya, A. M., Bullard, W. L., Kilbourne, J., Stern-Harbutte, A., Krajmalnik-Brown, R., Munk, B. H., Koppang, E. O., Lim, E. S., & Lucas, A. R. (2020). Immune protection is dependent on the gut microbiome in a lethal mouse gammaherpesviral infection. *Scientific Reports*, 10(1), 2371. <https://doi.org/10.1038/s41598-020-59269-9>
- [6] Walls, A. C., Park, Y. J., Tortorici, M. A., Wall, A., McGuire, A. T., & Veasley, D. (2020). Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell*, 181(2), 281–92.e6.
- [7] Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., ... Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*, 395(10223), 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- [8] Mahajan, A., & Hirsch, J. A. (2020). Novel coronavirus: What neuroradiologists as citizens of the world need to know. *AJNR. American Journal of Neuroradiology*, 41(4), 552–554. <https://doi.org/10.3174/ajnr.A6526>
- [9] WHO. (2020). COVID-19: Operational guidance for maintaining essential health services during an outbreak: Interim guidance, 25 March 2020. WHO.
- [10] Lai, C. C., Shih, T. P., Ko, W. C., Tang, H. J., & Hsueh, P. R. (2020). Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019

- (COVID-19): The epidemic and the challenges. *International Journal of Antimicrobial Agents*, 55(3), 105924. <https://doi.org/10.1016/j.ijantimicag.2020.105924>
- [11] Poyiadji, N., Shahin, G., Noujaim, D., Stone, M., Patel, S., & Griffith, B. (2020). COVID-19–associated acute hemorrhagic necrotizing encephalopathy: Imaging features. *Radiology*, 296(2), E119–E120. <https://doi.org/10.1148/radiol.2020201187>
- [12] Mao, L., Jin, H., Wang, M., Hu, Y., Chen, S., He, Q., Chang, J., Hong, C., Zhou, Y., Wang, D., Miao, X., Li, Y., & Hu, B. (2020). Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurology*, 77(6), 683–690. <https://doi.org/10.1001/jamaneurol.2020.1127>
- [13] Novel, C. P. E. R. E., & the Epidemiology Working Group for NCIP Epidemic Response, Chinese Center for Disease Control and Prevention. (2020). [The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 41(2), 145–151.
- [14] Maddah, S. A., & Modanloo, M. (2020). The challenges of keeping psychiatric patients safe in rehabilitation centers during coronavirus outbreak. *Archives of Psychiatric Nursing*, 34(6), 580–581. <https://doi.org/10.1016/j.apnu.2020.07.024>
- [15] Noé, A., Ribeiro, R. M., Anselmo, R., Maixenchs, M., Sitole, L., Munguambe, K., Blanco, S., le Souef, P., & García-Basteiro, A. L. (2017). Knowledge, attitudes and practices regarding tuberculosis care among health workers in Southern Mozambique. *BMC Pulmonary Medicine*, 17(1), 2. <https://doi.org/10.1186/s12890-016-0344-8>
- [16] Dauda Goni, M., Hasan, H., Naing, N. N., Wan-Arfah, N., Zeiny Deris, Z., Nor Arifin, W., & Abubakar Baaba, A. (2019). Assessment of knowledge, attitude and practice towards prevention of respiratory tract infections among Hajj and Umrah Pilgrims from Malaysia in 2018. *International Journal of Environmental Research and Public Health*, 16(22), 4569. <https://doi.org/10.3390/ijerph16224569>
- [17] Larson, E. L., Bryan, J. L., Adler, L. M., & Blane, C. (1997). A multifaceted approach to changing handwashing behavior. *American Journal of Infection Control*, 25(1), 3–10. [https://doi.org/10.1016/S0196-6553\(97\)90046-8](https://doi.org/10.1016/S0196-6553(97)90046-8)
- [18] Goni, M. D., Naing, N. N., Hasan, H., Wan-Arfah, N., Deris, Z. Z., Arifin, W. N., Hussin, T. M. A. R., Abdulrahman, A. S., Baaba, A. A., & Arshad, M. R. (2020). Development and validation of knowledge, attitude and practice questionnaire for prevention of respiratory tract infections among Malaysian Hajj pilgrims. *BMC Public Health*, 20(1), 189. <https://doi.org/10.1186/s12889-020-8269-9>
- [19] Ayed, A., & Zabn, K. (2021). Knowledge and attitude towards COVID-19 among nursing students: Palestinian perspective. *SAGE Open Nursing*, 7(7), 23779608211015150. <https://doi.org/10.1177/23779608211015150>

- [20] Quisao, E. Z. S., Tayaba, R. R. R., & Soriano, G. P. (2021). Knowledge, attitude, and practice towards COVID-19 among student nurses in Manila, Philippines: A cross-sectional study. *Belitung Nursing Journal*, 7(3), 203–209. <https://doi.org/10.33546/bnj.1405>
- [21] Le An, P., Huynh, G., Nguyen, H. T. N., Pham, B. D. U., Nguyen, T. V., Tran, T. T. T., & Tran, T. D. (2021). Knowledge, attitude, and practice towards COVID-19 among healthcare students in Vietnam. *Infection and Drug Resistance*, 14, 3405–3413. <https://doi.org/10.2147/IDR.S328677>
- [22] Sahin, M. K., Aker, S., & Kaynar Tuncel, E. (2015). Knowledge, attitudes and practices concerning Middle East respiratory syndrome among Umrah and Hajj pilgrims in Samsun, Turkey, 2015. *Eurosurveillance*, 20(38), 30023. <https://doi.org/10.2807/1560-7917.ES.2015.20.38.30023>
- [23] Sridhar, S., Belhouchat, K., Drali, T., Benkouiten, S., Parola, P., Brouqui, P., & Gautret, P. (2015). French Hajj pilgrims' experience with pneumococcal infection and vaccination: A knowledge, attitudes and practice (KAP) evaluation. *Travel Medicine and Infectious Disease*, 13(3), 251–255. <https://doi.org/10.1016/j.tmaid.2015.02.002>
- [24] Gautret, P., Soula, G., Parola, P., & Brouqui, P. (2009). Hajj pilgrims' knowledge about acute respiratory infections. *Emerging Infectious Diseases*, 15(11), 1861–1862. <https://doi.org/10.3201/eid1511.090201>
- [25] Alhomoud, F., & Alhomoud, F. (2017). “Your health essential for your Hajj”: Muslim pilgrims' knowledge, attitudes and practices regarding Middle East respiratory syndrome coronavirus (MERS-CoV) during Hajj season. *Journal of Infection and Chemotherapy*, 23(5), 286–292. <https://doi.org/10.1016/j.jjiac.2017.01.006>
- [26] Asaad, A., El-Sokkary, R., Alzamanan, M., & El-Shafei, M. (2020). Knowledge and attitudes towards Middle East respiratory syndrome-coronavirus (MERS-CoV) among health care workers in south-western Saudi Arabia. *Eastern Mediterranean Health Journal*, 26(4), 435–442. <https://doi.org/10.26719/emhj.19.079>
- [27] Gautret, P., Benkouiten, S., Salaheddine, I., Belhouchat, K., Drali, T., Parola, P., & Brouqui, P. (2013). Hajj pilgrims knowledge about Middle East respiratory syndrome coronavirus, August to September 2013. *Eurosurveillance*, 18(41), 20604. <https://doi.org/10.2807/1560-7917.ES2013.18.41.20604>
- [28] Glanz, K., & Bishop, D. B. (2010). The role of behavioral science theory in development and implementation of public health interventions. *Annual Review of Public Health*, 31, 399–418. <https://doi.org/10.1146/annurev.publhealth.012809.103604>
- [29] Adli, I., Widyahening, I. S., Lazarus, G., Phowira, J., Baihaqi, L. A., Ariffandi, B., Putera, A. M., Nugraha, D., Gamalliel, N., & Findyartini, A. (2022). Knowledge,

attitude, and practice related to the COVID-19 pandemic among undergraduate medical students in Indonesia: A nationwide cross-sectional study. *PLoS One*, 17(1), e0262827. <https://doi.org/10.1371/journal.pone.0262827>

- [30] Farah, A. M., Nour, T. Y., Obsiye, M., Aden, M. A., Ali, O. M., Hussein, M. A., Budul, A. B., Omer, M., & Getnet, F. (2021). Knowledge, attitudes, and practices regarding COVID-19 among health care workers in public health facilities in Eastern Ethiopia: Cross-sectional Survey Study. *JMIR Formative Research*, 5(10), e26980. <https://doi.org/10.2196/26980>